

ORGANOMETALLIC COMPLEXES BEARING SALMEN AND SALBEN LIGANDS AND THEIR APPLICATION IN CATALYSIS

Stefano Baratti^{a,b}

^a Dipartimento di Scienze Chimiche e Geologiche, Università degli Studi di Modena e Reggio Emilia, via G. Campi 103, 41125 Modena, Italy

^b Department of Synthetic Molecules Technical Development, Process Chemistry & Catalysis, F. Hoffmann-La Roche Ltd, Basel, Switzerland

Contacts: stefano.baratti2@unimore.it

Abstract

H₂salmen and H₂salben (**Figure 1**) are Schiff bases formed by the condensation of salicylaldehyde (salH) with methylenediamine (men) and phenylenediamine (ben), respectively. These ligands are structurally related to the well-known H₂salen (employing ethylenediamine, en) and exhibit similar N₂O₂ coordination environment toward metal ions. However, due to the one-carbon chain between the iminic nitrogen atoms, H₂salmen and H₂salben tend to form dinuclear [1,2] or oligonuclear [3] metal complexes. These complexes typically adopt geometries that lie between square planar and tetrahedral [1,2], in contrast to the mononuclear square planar complexes, characteristic of H₂salen [4]. Copper [1], cobalt [2,3], iron [5,6], and manganese [5] readily form complexes with these ligands. The flexibility of the coordination environment and the stabilization of mixed oxidation state species play a crucial role in the development of efficient catalyst.




Figure 1: structure of H₂salmen, H₂salben and H₂salen ligands

Building on the successful application of copper(II)-salmen and salben complexes in C-N coupling reactions [7] as well as the established catalytic activity of metal-salen complexes [4], my PhD project aims to expand the understanding of metal-salmen/salben derivatives through a three-steps approach. First, efforts will focus on the synthesis of tuned salmen and salben ligands, building upon the expertise developed in our research group, followed by complexation with first-row metal ions, particularly cobalt [2]. Next, the catalytic properties of the new obtained complexes will be evaluated in selected benchmark, focusing on the performances in coupling reaction, including C-C, C-O and C-N bond formation. Finally, mechanistic investigation will be carried out to gain deeper insight into the catalytic pathways, enabling the rational design of next-generation metal complexes.

References

- [1] Alessandro Pasini et al., *J. Chem. Soc., Dalton Trans.*, **2000**, 3467–3472.
- [2] Brunetto Chiari et al., *J. Chem. Soc., Dalton Trans.*, **2001**, 3611–3616.
- [3] Brunetto Chiari et al., *J. Chem. Soc., Dalton Trans.*, **2002**, 4672–4677.
- [4] Pier Giorgio Cozzi, *Chem. Soc. Rev.*, **2004**, 33, 410–421.
- [5] Luca Rigamonti et al., *Int. J. Mol. Sci.*, **2020**, 21, 7882.
- [6] Lorenzo Marchi et al. *Int. J. Mol. Sci.*, **2023**, 24, 5808.
- [7] Serena Maria Fantasia et al., Patent, **2025**, WO 2025/016888 Al.